



ZYMO RESEARCH

The Beauty of Science is to Make Things Simple

INSTRUCTION MANUAL

ZR Plasmid Miniprep™-Classic

Catalog Nos. **D4015, D4016, & D4054**

Highlights

- For purification of high quality, endotoxin-free plasmid DNA for restriction endonuclease digestion, DNA sequencing, transformation, cloning, transfection, *in vitro* transcription reactions, etc.
- Innovative colored buffers* for rapid identification of complete bacterial cell lysis and neutralization steps.
- Unique column design: zero buffer retention and low volume (30 µl) elution.

Contents

Product Contents.....	1
Specifications	1
Product Description	2
Buffer Preparation.....	3
Protocol	3
Troubleshooting.....	4, 5
Ordering Information.....	6
Other Products	7

* Patent Pending; For Research Use Only

Ver. 1.0.4

ZYMO RESEARCH CORP.

Toll Free: 1-888-882-9682 • Fax: • Web: www.zymoresearch.com • E-mail: info@zymoresearch.com

Satisfaction of all Zymo Research products is guaranteed. If you are not satisfied with this product please call 1-888-882-9682.

Product Contents

ZR Plasmid Miniprep™-Classic (Kit Size)	D4015 (100 preps.)	D4016 (400 preps.)	D4054 (800 preps.)	Storage Temperature
P1 Buffer (Red)	20 ml	80 ml	160 ml	Room Temp.
P2 Buffer ¹ (Green)	20 ml	80 ml	160 ml	Room Temp.
P3 Buffer ² (Yellow)	50 ml	220 ml	2 x 220 ml	4° after opening.
Endo-Wash Buffer	30 ml	2 x 60 ml	3 x 60 ml	Room Temp.
Plasmid Wash Buffer (concentrate) ²	24 ml	48 ml	2 x 48 ml	Room Temp.
DNA Elution Buffer	4 ml	16 ml	2 x 16 ml	Room Temp.
Zymo-Spin™ IIN Columns	100	400	800	Room Temp.
Collection Tubes	100	400	800	Room Temp.
Instruction Manual	1	1	1	-

Note: Integrity of kit components is guaranteed for up to one year from date of purchase. Reagents are routinely tested on a lot-to-lot basis to ensure they provide maximal performance and reliability.

¹ Caution: **P2 Buffer** contains NaOH and **P3 Buffer** contains chaotropic reagents. Please use proper safety precautions with these reagents.

² Add ethanol to **Plasmid Wash Buffer** (concentrate) prior to use. See **Buffer Preparation** (page 3) for instructions.

Specifications:

- **DNA Purity:** High purity, endotoxin-free (<50 EU/μg) plasmid DNA eluted in low salt buffer or water; typical $A_{(260/280)} \geq 1.8$. DNA is suitable for restriction endonuclease digestion, sequencing, transfection, ligation, *in vitro* transcription, labeling, and other reactions requiring highly purified DNA.
- **Recovery Volume:** ≥30 μl
- **Plasmid DNA Size:** Up to 25 kb
- **Plasmid DNA yield:** Up to 25 μg per preparation depending on the plasmid copy number, input volume of *E. coli* culture, and culture growth conditions.
- **Procedure:** Can be conducted at room temperature, between 15 - 30°C.

Note: ™ Trademarks of Zymo Research Corporation. This product is for research use only and should only be used by trained professionals. It is not intended for use in diagnostic procedures. Some reagents included with this kit are irritants. Wear protective gloves and eye protection. Follow the safety guidelines and rules enacted by your research institution or facility.

Several ZR Plasmid Miniprep™-Classic product technologies are subject to U.S. and foreign patents or are patent pending.

ZYMO RESEARCH CORP.

Toll Free: 1-888-882-9682 • Fax: 1-949-266-9452 • Web: www.zymoresearch.com • E-mail: info@zymoresearch.com


Product Description

The **ZR Plasmid Miniprep™-Classic** kit is designed for efficient isolation of plasmid DNA from *E. coli* cell lysates using a procedure that is simple, rapid, user-friendly, and reliable compared to the products offered by the competition. It features a modified alkaline lysis protocol together with a unique *Fast Spin* column to yield high quality plasmid DNA in minutes. The ZR Plasmid Miniprep™-Classic features color-coded (red, green, yellow) reagents for easy determination of complete cell lysis. The innovative **Zymo-Spin™ IIN** columns facilitate high yield plasmid DNA that is endotoxin-free. Plasmid DNA purified using the ZR Plasmid Miniprep™-Classic kit is well suited for use in restriction endonuclease digestion, sequencing, DNA ligation, cloning, PCR, bacterial transformation, transfection, etc.


For **Technical Assistance**, please contact those at **Zymo Research's Technical Department** at 1-888-882-9682 or E-mail tech@zymoresearch.com.



Visualize complete bacterial cell lysis with unique colored **P1**, **P2**, and **P3** buffers.



Endonuclease digestion of three different DNA plasmids prepared using the **ZR Plasmid Miniprep™-Classic** (performed in triplicate). **M**: ZR 1 kb DNA Marker.



DNA sequencing chromatogram of plasmid DNA prepared using the **ZR Plasmid Miniprep™-Classic**.



The Zymo-Spin™ IIN Column: binds up to 25 µg DNA, 900 µl capacity, and 30 µl elution volume.

Buffer Preparation:

1. Add ethanol to the **Plasmid Wash Buffer** at a 4:1 volume ratio of ethanol to buffer.
 - For product **D4015**, add 96 ml 95 % ethanol to 24 ml **Plasmid Wash Buffer**.
 - For product **D4016**, add 192 ml 95 % ethanol to 48 ml **Plasmid Wash Buffer**.
 - For product **D4054**, add 192 ml 95 % ethanol to each 48 ml **Plasmid Wash Buffer**.

Notes:

¹The following procedures are carried out at a room temperature. All centrifugation steps should be performed between 11,000 - 16,000 x g.

²Depending on the volume of bacterial culture it may be necessary to repeat Step 1 several times.

³Excessive lysis can result in denatured plasmid DNA formation. When processing a large number of samples, work with groups of ≤ 10 at a time.

⁴A green precipitate consisting of K₂SDS and cell debris will form. A good way to mix is to shake the tube gently several times while it is inverted.

⁵The capacity of the collection tube with the column inserted is 800 µl. Empty the collection tube whenever necessary to prevent contamination of the spin column with the flow-through.

⁶Ensure that ethanol has been added to the concentrated **Plasmid Wash Buffer** prior to use.

⁷The **DNA Elution Buffer** contains 10 mM Tris-HCl, pH 8.5, 0.1 mM EDTA. If required, pure water can be used to elute the DNA. Add the **DNA Elution Buffer** directly to the center of the **Zymo-Spin™ IIN** column matrix to ensure optimal DNA elution.

Protocol¹


1. Centrifuge 0.5 - 5 ml^{1,2} of bacterial culture in a clear 1.5 ml tube at full speed for 15 - 20 seconds in a microcentrifuge. Discard supernatant.
2. Add 200 µl of **P1 Buffer** (Red) to the tube and resuspend pellet completely (i.e., by vortexing or pipeting).
3. Add 200 µl of **P2 Buffer** (Green)³ and mix by inverting the tube 2 - 4 times. Cells are completely lysed when the solution appears clear, purple, and viscous. Proceed to the next step within 1-2 minutes.
4. Add 400 µl of **P3 Buffer** (Yellow) and mix gently but thoroughly. Do not vortex. The sample will turn yellow when the neutralization is complete⁴. Allow the lysate to incubate at room temperature for 1-2 minutes.
5. Centrifuge sample(s) for 2 minutes.
6. Place a **Zymo-Spin™ IIN** column in a **Collection Tube** and transfer the supernatant from Step 5 into the **Zymo-Spin™ IIN** column. When pipeting the supernatant be careful not to disturb the green pellet to avoid transferring any cellular debris to the column.
7. Centrifuge the **Zymo-Spin™ IIN/Collection Tube** assembly for 30 seconds.
8. Discard the flow-through in the **Collection Tube**, making sure the flow-through does not touch the bottom of the column. Return the **Zymo-Spin™ IIN** column to the **Collection Tube**⁵.
9. Add 200 µl of **Endo-Wash Buffer** to the column and centrifuge for 30 seconds.
10. Add 400 µl of **Plasmid Wash Buffer**⁶ to the column. Centrifuge for 1 minute.
11. Transfer the column into a clean 1.5 ml microcentrifuge tube and then add 30 µl (of **DNA Elution Buffer**⁷ to the column. Centrifuge for 30 seconds to elute the plasmid DNA.

Troubleshooting Guide:

Problem	Possible Causes and Suggested Solutions
Low DNA Yield	

Culture growth conditions

- Poor aeration of culture. The optimal culture volume to air volume ratio is 1:4 or less (20% culture, 80% air). For best aeration, use baffled culture flasks, a vented or gas-permeable seal on the culture vessel, and incubate with vigorous shaking.



- Other Possible reasons may include: An overgrown/undergrown or contaminated culture, or omission of antibiotics from the growth medium. Use a fresh culture for optimal performance. Grow the culture to an O.D.₆₀₀ >1.0.

Procedural errors


- Incomplete lysis: After addition of P2 Buffer the solution should change from opaque red to clear purple, indicating complete lysis. Different *E. coli* strains often require different growth conditions and may vary in their susceptibility to alkaline lysis.
- Incomplete neutralization: Cell debris will float to the surface after centrifugation and the pellet may appear “puffy”. Make sure the neutralization is complete prior to centrifugation. Invert the tube an additional 2 - 3 times after the sample turns yellow following the addition of P3 Buffer.

Plasmid Wash Buffer

- Ensure that ethanol has been added to the wash buffer.

DNA elution

- Incomplete elution: For large size plasmids (> 10 kb), incubate the column for 5 - 10 minutes before centrifugation. Also, pre-warm the DNA Elution Buffer to 50 °C prior to elution and increase the elution volume to $\geq 50 \mu\text{l}$.

**Low DNA Quality***DNA does not perform well*

- Incomplete neutralization generates poor quality supernatant and results in loading too much cell debris onto the column. Ensure that neutralization is complete by inverting the sample an additional 2 - 3 times after the addition of P3 Buffer.
- The spin column tip is contaminated with wash buffer flowthrough. Avoid tilting the collection tube after the last wash step to ensure that the column tip does not contact the flowthrough. Empty the collection tube when recommended in the protocol.
- Insufficient centrifugation: make sure that all centrifugation steps are performed between 11,000 - 16,000 x g. If a lower centrifuge speed is used, then extend the centrifugation time to compensate.

RNA in eluate

- After neutralization, be sure to allow lysate to incubate 1-2 minutes before centrifugation.

Genomic DNA in eluate

- Improper handling (sample was vortexed or handled too roughly after the addition of P2 & P3 Buffer). Genomic DNA contamination is usually caused by excessive mechanical shearing during the lysis and neutralization steps. Also, prolonged lysis or incomplete mixing of lysis or neutralization buffers may contribute to genomic DNA contamination in your sample.
- Overgrown culture. Older cultures may contain more genomic DNA contamination than fresh cultures.

Ordering Information

Product Description	Kit Size	Catalog No.
ZR Plasmid Miniprep™-Classic	100 preps.	D4015
	400 preps.	D4016
	800 preps.	D4054

For Individual Sale	Amount	Catalog No.
P1 Buffer (Red)	20 ml	D4027-1-20
	80 ml	D4027-1-80
	160 ml	D4027-1-160
P2 Buffer (Green)	20 ml	D4027-2-20
	80 ml	D4027-2-80
	160 ml	D4027-2-160
P3 Buffer (Yellow)	50 ml	D4027-3-50
	220 ml	D4027-3-220
	440 ml	D4027-3-440
Endo-Wash Buffer	30 ml	D4036-3-30
	60 ml	D4036-3-60
Plasmid Wash Buffer (concentrate)	24 ml	D4027-4-24
	48 ml	D4027-4-48
DNA Elution Buffer	4 ml	D3004-4-4
	10 ml	D3004-4-10
	16 ml	D3004-4-16
Zymo-Spin IIN™ Columns	50 columns	C1019-50
	250 columns	C1019-250
Collection Tubes	50 tubes	C1001-50
	500 tubes	C1001-500
	1000 tubes	C1001-1000

Other Popular DNA Purification Products from Zymo Research

Product	Format	Kit Size	Cat No.
Fragment DNA Clean-up, Concentration & Recovery			
DNA Clean & Concentrator™-5	Spin Column Format (up to 5 µg/prep)	50 preps. 200 preps.	D4003*, D4013 D4004*, D4014
DNA Clean & Concentrator™-25	Spin Column Format (up to 25 µg/prep)	50 preps. 200 preps.	D4005*, D4033 D4006*, D4034
DNA Clean & Concentrator™-100	Spin Column Format (up to 100 µg/prep)	25 preps. 50 preps.	D4029 D4030
DNA Clean & Concentrator™-500	Spin Column Format (up to 500 µg/prep)	10 preps. 20 preps.	D4031 D4032
ZR-96 DNA Clean & Concentrator™-5	96-Well Format (up to 5 µg/well; deep well)	2 x 96 preps. 4 x 96 preps.	D4023 D4024
ZR-96 DNA Clean-up Kit™	96-Well Format (up to 5 µg/well; shallow well)	2 x 96 preps. 4 x 96 preps.	D4017 D4018
ZR DNA Sequencing Clean-up Kit™	Spin Column Format (up to 5 µg/prep)	50 preps. 200 preps.	D4050 D4051
ZR-96 DNA Sequencing Clean-up Kit™	96-Well Format (up to 5 µg/well)	2 x 96 preps. 4 x 96 preps.	D4052 D4053
OneStep™ PCR Inhibitor Removal Kit	Spin Column Format (up to 25 µg/prep)	50 preps.	D6030
OneStep-96™ PCR Inhibitor Removal Kit	96-Well Format (up to 5 µg/well)	2x96 preps.	D6035
Zymoclean™ Gel DNA Recovery Kit	Spin Column Format (up to 5 µg/prep)	50 preps. 200 preps.	D4001 D4002
ZR-96 Zymoclean™ Gel DNA Recovery Kit	96-Well Format (up to 5 µg/well)	2 x 96 preps. 4 x 96 preps.	D4021 D4022
Plasmid DNA Isolation			
Zyppy™ Plasmid Miniprep Kit	Pellet Free, Spin Column Format	50 preps. 100 preps. 400 preps. 800 preps.	D4036 D4019 D4020 D4037
Zyppy™ Plasmid Midiprep Kit	Pellet Free, Spin Column Format	25 preps. 50 preps.	D4025 D4026
Zyppy™ Plasmid Maxiprep Kit	Spin/Vacuum Column Format	10 preps. 20 preps.	D4027 D4028
Genomic DNA Isolation			
ZR Genomic DNA I Kit™	Silica Bead Format - Scaleable	100 preps. 400 preps.	D3004 D3005
ZR Genomic DNA II Kit™	Spin Column Format (up to 25 µg/prep)	50 preps. 200 preps.	D3006*, D3024 D3007*, D3025
ZR-96 Genomic DNA Kit™	96-Well Format (up to 5 µg/well)	2 x 96 preps. 4 x 96 preps. 10 x 96 preps.	D3010 D3011 D3012
ZR Genomic DNA™-Tissue MiniPrep	Spin Column Format (up to 25 µg/prep)	50 preps. 200 preps.	D3050 D3051
ZR-96 Genomic DNA™-Tissue MiniPrep	96-Well Format (up to 5 µg/well)	2 x 96 preps. 4 x 96 preps. 10 x 96 preps.	D3055 D3056 D3057
Pinpoint™ Slide DNA Isolation System	For Archived Tissue Sections, Spin Column Format (up to 5 µg/prep)	50 preps.	D3001
ZR Serum DNA Kit™	Silica Bead Format - Scaleable	scaleable	D3013
ZR Urine DNA Isolation Kit™	Filtration, Spin Column Format (up to 5 µg/prep)	20 preps.	D3060
ZR Viral DNA Kit™	Spin Column Format (up to 5 µg/prep)	50 preps. 200 preps.	D3015 D3016
ZR-96 Viral DNA Kit	96-Well Format (up to 5 µg/well)	2 x 96 preps. 4 x 96 preps.	D3017 D3018
Environmental DNA Isolation			
ZR Soil Microbe DNA Kit™	Bead Bashing, Spin Column Format (up to 25 µg/prep)	50 preps.	D6001
ZR-96 Soil Microbe DNA Kit™	Bead Bashing, 96-Well Format (up to 5 µg/well)	2 x 96 preps.	D6002
ZR Fungal/Bacterial DNA Kit™	Bead Bashing, Spin Column Format (up to 25 µg/prep)	50 preps.	D6005
ZR-96 Fungal/Bacterial DNA Kit™	Bead Bashing, 96-Well Format (up to 5 µg/well)	2 x 96 preps.	D6006
ZR Fecal DNA Kit™	Bead Bashing, Spin Column Format (up to 25 µg/prep)	50 preps.	D6010
ZR-96 Fecal DNA Kit™	Bead Bashing, 96-Well Format (up to 5 µg/well)	2 x 96 preps.	D6011
ZR Tissue & Insect DNA Kit-5™	Bead Bashing, Spin Column Format (up to 5 µg/prep)	50 preps.	D6015
ZR Tissue & Insect DNA Kit-25™	Bead Bashing, Spin Column Format (up to 25 µg/prep)	50 preps.	D6016
ZR-96 Tissue & Insect DNA Kit™	Bead Bashing, 96-Well Format (up to 5 µg/well)	2 x 96 preps.	D6017
ZR Plant/Seed DNA Kit™	Bead Bashing, Spin Column Format (up to 25 µg/prep)	50 preps.	D6020
ZR-96 Plant/Seed DNA Kit™	Bead Bashing, 96-Well Format (up to 5 µg/well)	2 x 96 preps.	D6021

* Uncapped Spin Column Format

ZYMO RESEARCH CORP.

Toll Free: 1-888-882-9682 • Fax: 1-949-266-9452 • Web: www.zymoresearch.com • E-mail: info@zymoresearch.com